

Lagrangian fibrations in four and six dimensions

Justin Sawon¹



THE UNIVERSITY
of **NORTH CAROLINA**
at **CHAPEL HILL**

Algebraic Geometry Seminar
Milano, 3 July, 2023

¹Supported by MPIM Bonn and NSF awards [DMS-1555206](#), [DMS-2152130](#).

Overview

- generalities on Lagrangian fibrations
- examples in four dimensions
- examples in six dimensions
- classification results
- polarization types

(partly joint work with Chen Shen and Xuqiang Qin)

Holomorphic symplectic manifolds

Let X be a compact Kähler manifold with $c_1 = 0$.

Thm (Bogomolov): \exists finite étale cover \tilde{X} of X with

$$\tilde{X} = T \times \prod_i CY_i \times \prod_j IHS_j,$$

$T =$ torus, $CY_i =$ (strict) Calabi-Yau manifolds, and $IHS_j = \dots$

Def: A compact Kähler manifold X is a *holomorphic symplectic manifold* if it admits a non-degenerate holomorphic two-form σ .

In addition if $\pi_1(X) = 0$ and $H^0(\Omega^2)$ is generated by σ then we say X is an *irreducible holomorphic symplectic (IHS) manifold*.

Examples of IHS manifolds

1. Hilbert schemes of points on K3 surfaces, $\text{Hilb}^n S \rightarrow \text{Sym}^n S$.
2. Generalized Kummer varieties, $\widetilde{\text{Hilb}}^{n+1} A = A \times K_n(A)$.
Equivalently $K_n(A) := \text{kernel}(\text{Hilb}^{n+1} A \rightarrow \text{Sym}^{n+1} A \rightarrow A)$.
3. Mukai moduli spaces of stable sheaves on K3/abelian surfaces.

$$\text{Ext}^1(\mathcal{E}, \mathcal{E}) \times \text{Ext}^1(\mathcal{E}, \mathcal{E}) \rightarrow \text{Ext}^2(\mathcal{E}, \mathcal{E}) \xrightarrow{\text{tr}} \text{H}^2(\mathcal{O}) \cong \mathbb{C}$$

4. O'Grady's spaces, OG6 and OG10.

Up to deformation, two/three (smooth) examples known in dimensions four/six: $\text{Hilb}^2 S$, $K_2(A)$, $\text{Hilb}^3 S$, $K_3(A)$, and OG6.

Fibrations

Let X be an IHS manifold of dimension $2n$.

Thm (Matsushita): If $X \rightarrow B$ is a proper fibration then

1. $\dim B = n = \dim F$,
2. F is Lagrangian wrt the holomorphic symplectic form σ ,
3. generic fibre is a complex torus.

Thm (Hwang): B is isomorphic to \mathbb{P}^n if it is smooth.

Thm (Huybrechts-Xu): B is smooth if $n = 2$, thus $B \cong \mathbb{P}^2$.

Rmk (Voisin): Hodge theory \implies general fibre is an abelian variety.

Examples for $\text{Hilb}^n S$

1a. Beauville-Mukai system: Let C be a genus n curve in a K3 surface S , with $|C| \cong \mathbb{P}^n$ and \mathcal{C}/\mathbb{P}^n the family of curves linearly equivalent to C .

$$X := \overline{\text{Jac}}^d(\mathcal{C}/\mathbb{P}^n) \longrightarrow \mathbb{P}^n$$

is a Lagrangian fibration, deformation equivalent to $\text{Hilb}^n S$.

Or $X \cong$ moduli space $M(0, [C], 1 - g + d)$ of stable sheaves on S .

1b. If $S \longrightarrow \mathbb{P}^1$ is an elliptic K3 surface then the Hilbert scheme

$$\text{Hilb}^n S \rightarrow \text{Sym}^n S \rightarrow \text{Sym}^n \mathbb{P}^1 = \mathbb{P}^n$$

is a Lagrangian fibration. Its fibres look like

$$E_1 \times E_2 \times \cdots \times E_n.$$

(Generalized) Prym varieties

Let $\pi : C \rightarrow D$ be a double cover of curves with covering involution τ . Then

$$\text{Fix}^0(\tau^*) = \pi^* \text{Jac}^0 D \subset \text{Jac}^0 C.$$

Def: The Prym variety of C/D is an abelian variety

$$\text{Prym}(C/D) := \text{Fix}^0(-\tau^*),$$

of dimension $g_C - g_D$, principally polarized if $\pi : C \rightarrow D$ has zero or two branch points, otherwise polarization type

$$\underbrace{(1, \dots, 1)}_{g_C - 2g_D}, \underbrace{(2, \dots, 2)}_{g_D}.$$

Families of Prym varieties

Let $\pi : S \rightarrow T$ be a K3 double cover of another surface with *anti-symplectic* covering involution τ .

Thm (Nikulin): There exist 75 anti-symplectic involutions τ on K3s. The quotient $T = S/\tau$ is an Enriques or a rational surface.

A curve $D \subset T$ has a double cover $C \subset S$,

$$\begin{array}{ccc} C & \subset & S \\ 2:1 \downarrow & & 2:1 \downarrow \\ D & \subset & T. \end{array}$$

Let $\mathcal{D} \rightarrow |D|$ be the complete linear system in T , let $\tilde{\mathcal{C}} \rightarrow |C|$ be the complete linear system in S , and let

$$\mathcal{C} := \pi^* \mathcal{D} \subset \tilde{\mathcal{C}}.$$

Families of Prym varieties

There are two commuting *anti-symplectic* involutions on the Beauville-Mukai system $\overline{\text{Jac}}^0(\tilde{\mathcal{C}}/|C|)$:

- the involution τ^* induced by τ ,
- fibrewise duality $\mathcal{E} \mapsto \mathcal{E}xt_{\mathcal{S}}^1(\mathcal{E}, \mathcal{O}(-C))$ (takes $\iota_*\mathcal{L} \mapsto \iota_*\mathcal{L}^\vee$).

Thm (Markushevich-Tikhomirov, Arbarello-Saccà-Ferretti, Matteini): We can construct a relative Prym variety

$$\text{Prym}(\mathcal{C}/\mathcal{D}) := \text{Fix}^0(\mathcal{E} \mapsto \mathcal{E}xt_{\mathcal{S}}^1(\tau^*\mathcal{E}, \mathcal{O}(-C))) \subset \overline{\text{Jac}}^0(\tilde{\mathcal{C}}/|C|).$$

This is a symplectic *variety* and a Lagrangian fibration over $|D|$.

(1, 2)-polarized examples

3a. Markushevich-Tikhomirov system: S/T a K3 double cover of a degree two del Pezzo, C/D a genus three cover of an elliptic curve, $\text{Prym}(C/D)$ an abelian surface of type (1, 2).

Then $\text{Prym}(C/D) \rightarrow \mathbb{P}^2$ is an *irreducible* symplectic orbifold of dimension four, with 28 isolated $\mathbb{C}^4/\pm 1$ singularities.

Rmk: This orbifold is a partial resolution of the quotient of $\text{Hilb}^2 S$ by a symplectic involution, sometimes called the *Nikulin variety*.

(1, 2)-polarized examples

Another fibration on the Nikulin variety is constructed as follows.

3b. $F[2]$ acts fibrewise by translation on the Kummer K3

$$S \rightarrow E \times F / \pm 1 \rightarrow E / \pm 1 \cong \mathbb{P}^1$$

and there is an induced fibrewise action on $\text{Hilb}^2 S \rightarrow \mathbb{P}^2$.

Rmk: Each element of $F[2]$ acts as a symplectic involution.

Quotient by the action of a single element and blow-up the K3 of singularities to get an orbifold X .

Prop: X is an isotrivial Lagrangian fibration over \mathbb{P}^2 .

Rmk: Fibres $F \times F / \mathbb{Z}_2$ are (1, 2)-polarized. Moreover, X has $b_2 = 16$, $b_3 = 0$, $b_4 = 178$, and 28 isolated $\mathbb{C}^4 / \pm 1$ singularities.

Principally polarized examples

2a. Arbarello-Saccà-Ferretti system: S/T a K3 double cover of an Enriques, D genus $n + 1$, $\text{Prym}(C/D)$ principally polarized.

Then $\text{Prym}(C/D) \rightarrow \mathbb{P}^n$ is a symplectic variety, which is

- birational to a Beauville-Mukai system if D is hyperelliptic,
- simply connected with no symplectic resolution otherwise,
- and irreducible if n is even.

If $n = 2$ or 3 it has isolated $\mathbb{C}^4 / \pm 1$ or $\mathbb{C}^6 / \pm 1$ singularities.

Lemma: If $C = C_1 \cup C_2$ with $C_1.C_2 = 2k$ then a neighbourhood of $[\mathcal{F}_1 \oplus \mathcal{F}_2] \in \text{Prym}(C/D)$ looks locally like $\mathbb{C}^{N-2k} \times (\mathbb{C}^{2k} / \pm 1)$.

Principally polarized examples

2b. Let S be a Kummer K3 surface with an elliptic fibration

$$S \longrightarrow E \times F / \pm 1 \longrightarrow E / \pm 1 \cong \mathbb{P}^1.$$

$\mathrm{Hilb}^2 S \rightarrow \mathbb{P}^2$ is an isotrivial fibration with smooth fibres $F \times F$.

The group $F[2] \cong \mathbb{Z}_2^{\oplus 2}$ acts by diagonal translation on $F \times F$ and fibrewise on $\mathrm{Hilb}^2 S$. Take the quotient $\mathrm{Hilb}^2 S / \mathbb{Z}_2^{\oplus 2}$ and blow-up codimension two singularities to get a symplectic orbifold X .

Prop: X is an isotrivial Lagrangian fibration over \mathbb{P}^2 .

Rmk: Fibres $F \times F / F[2]$ are principally polarized. Moreover, X has $b_2 = 14$, $b_3 = 0$, $b_4 = 150$, and 36 isolated $\mathbb{C}^4 / \pm 1$ singularities.

Examples for $K_n(A)$

4a. Debarre system: Let $C \subset A$ give a polarization of type $(1, n+1)$. Then C has genus $n+2$ and $|C| \cong \mathbb{P}^n$. Let \mathcal{C}/\mathbb{P}^n be the family of curves linearly equivalent to C and

$$Y := \overline{\text{Jac}}^d(\mathcal{C}/\mathbb{P}^n) \longrightarrow \mathbb{P}^n.$$

We get a Lagrangian fibration

$$X := \text{kernel}(\text{Alb} : Y \longrightarrow A)$$

with $(1, \dots, 1, n+1)$ -polarized fibres $X_t = \ker(\overline{\text{Jac}}^d C_t \longrightarrow A)$.

4b. If $A = E \times F$ then

$$\text{Hilb}^{n+1}A \longrightarrow \text{Sym}^{n+1}A \longrightarrow \text{Sym}^{n+1}E \longrightarrow J^{n+1}E \cong E$$

induces an isotrivial Lagrangian fibration $K_n(A) \longrightarrow \mathbb{P}^n$ with $(1, \dots, 1, n+1)$ -polarized fibres

$$\cong \{(f_0, f_1, \dots, f_n) \in F^{n+1} \mid f_0 + f_1 + \dots + f_n = 0 \text{ in } F\}.$$

Summary of examples in four dimensions

Example	Polarization type
Beauville-Mukai system	(1, 1)
Hilb ² <i>S</i> of an elliptic K3 <i>S</i>	(1, 1)
Arbarello-Saccà-Ferretti system	(1, 1)
Isotrivial system on Hilb ² <i>S</i> / $\mathbb{Z}_2^{\oplus 2}$	(1, 1)
Markushevich-Tikhomirov system	(1, 2)
Isotrivial system on Hilb ² <i>S</i> / \mathbb{Z}_2	(1, 2)
Debarre system for $A^{1,3}$	(1, 3)
Isotrivial system on $K_2(E \times F)$	(1, 3)

More six-dimensional examples

Matteini system: S/T a K3 double cover of a cubic del Pezzo, C/D a genus four cover of an elliptic curve, $\text{Prym}(C/D)$ an abelian threefold of type $(1, 1, 2)$.

$\text{Prym}(C/D) \rightarrow \mathbb{P}^3$ is an *irreducible* symplectic orbifold of \dim^n six, with singularities that look like $\mathbb{C}^2 \times (\mathbb{C}^4 / \pm 1)$ and $\mathbb{C}^6 / \mathbb{Z}_2 \times \mathbb{Z}_2$.

S-Shen system: S/T a K3 double cover of a degree one del Pezzo, $D \in |-2K_T|$, C/D a genus five cover of a genus two curve, $\text{Prym}(C/D)$ an abelian threefold of type $(1, 2, 2)$. Then


$$\text{Prym}(C/D) := \text{Fix}^0(-) \subset \overline{\text{Jac}}^0(\tilde{C}/|C|) \leftarrow \text{OG10}$$

is a symplectic variety of \dim^n six, with 120 isolated singularities that look like $\mathbb{C}^6 / \pm 1$.

A birational model

The del Pezzo T is a double cover of the quadric cone Q . The covering involution lifts to another anti-symplectic involution on S .

Their composition gives a symplectic involution on S , with quotient a singular K3 surface \bar{S} with 8 A_1 -singularities.



A generic τ -invariant $C \subset S$ is an étale double cover of a genus three curve $\bar{C} \subset \bar{S}$, which is isomorphic to $\tilde{C} \subset \tilde{S}$.

A birational model

Pull-back induces a map

$$\mathrm{Jac}^0 \tilde{\mathcal{C}} = \mathrm{Jac}^0 \overline{\mathcal{C}} \longrightarrow \mathrm{Jac}^0 \mathcal{C}$$

which is two-to-one onto its image $\mathrm{Prym}(\mathcal{C}/\mathcal{D})$.

Let $\tilde{\mathcal{M}} := \overline{\mathrm{Jac}^0}(\tilde{\mathcal{C}}/\mathbb{P}^3)$ be the Beauville-Mukai system of $\tilde{\mathcal{C}} \subset \tilde{\mathcal{S}}$. Then there is a rational dominant generically two-to-one map

$$\tilde{\mathcal{M}} \dashrightarrow \mathrm{Prym}(\mathcal{C}/\mathcal{D}).$$

Moreover, $\tilde{\mathcal{M}}$ is deformation equivalent to $\mathrm{Hilb}^3 \tilde{\mathcal{S}}$.

Thm (S-Shen): $\mathrm{Prym}(\mathcal{C}/\mathcal{D})$ is a *primitive* symplectic variety:

- the symplectic structure is unique up to a scalar, $h^{2,0} = 1$,
- we have vanishing of the Hodge number $h^{1,0} = 0$.

Examples for $K_3(A)$ and OG6

Debarre system/OG6: Let $C \subset A$ give a polarization of type $(2, 2)$. Then C has genus 5 and $|C| \cong \mathbb{P}^3$. Consider

$$Y := \overline{\text{Jac}}^d(C/\mathbb{P}^3) \longrightarrow \mathbb{P}^3,$$

i.e., $Y \cong M(0, [C], d - 4)$ on A , and $X := \text{kernel}(\text{Alb} : Y \longrightarrow A)$.

- If d is odd then X is deformation equivalent to $K_3(A)$.
- If d is even then \tilde{X} is deformation equivalent to OG6.

Rmk: Both cases have fibres of polarization type $(1, 2, 2)$.

Summary of examples in six dimensions

Example	Polarization type
Beauville-Mukai system	(1, 1, 1)
Hilb ³ S of an elliptic K3 S	(1, 1, 1)
Arbarello-Saccà-Ferretti system	(1, 1, 1)
Matteini system	(1, 1, 2)
S-Shen system	(1, 2, 2)
Debarre system for $A^{1,4}$	(1, 1, 4)
Isotrivial system on $K_3(A)$	(1, 1, 4)
Debarre system for $A^{2,2}$	(1, 2, 2)
O'Grady 6 on $A^{2,2}$	(1, 2, 2)

Fibrations by Jacobians

Thm (Markushevich): Let \mathcal{C}/\mathbb{P}^2 be a flat family of integral Gorenstein curves of genus two such that $X = \overline{\text{Jac}}^d(\mathcal{C}/\mathbb{P}^2)$ is a Lagrangian fibration (with X smooth!). Then $X \rightarrow \mathbb{P}^2$ must be a Beauville-Mukai integrable system.

Rmk: The general principally polarized abelian surface is the Jacobian of a genus two curve.

Thm (Matsushita): $R^i \pi_* \mathcal{O}_X \cong \Omega_{\mathbb{P}^2}^i$.

When $i = 1$ this says $TF \cong N_{F/CX}^\vee$ for smooth fibres F .

Moreover, we have $R^1 \pi_* \mathcal{O}_{\mathcal{C}} \cong R^1 \pi_* \mathcal{O}_X \cong \Omega_{\mathbb{P}^2}^1$.

Fibrations by Jacobians

Proof: The relative canonical map gives a double cover

$$\mathcal{C} \longrightarrow \mathbb{P}(R^1\pi_*\mathcal{O}_{\mathcal{C}}) = \mathbb{P}(R^1\pi_*\mathcal{O}_X) = \mathbb{P}(\Omega_{\mathbb{P}^2}^1) \subset \mathbb{P}^2 \times (\mathbb{P}^2)^\vee$$

branched over the zero locus in $\mathbb{P}(\Omega_{\mathbb{P}^2}^1)$ of a section of

$$\mathcal{O}_{\mathbb{P}(\Omega^1)}(6) \otimes \pi^*\mathcal{O}_{\mathbb{P}^2}(2k) = \mathcal{O}(2k + 6, 6)|_{\mathbb{P}(\Omega^1)}.$$

Now $R^1\pi_*\mathcal{O}_{\mathcal{C}} = \Omega_{\mathbb{P}^2}^1$ determines $k = -3$, so

$$\mathcal{O}(2k + 6, 6)|_{\mathbb{P}(\Omega^1)} = \mathcal{O}(0, 6)|_{\mathbb{P}(\Omega^1)}$$

is pulled back from $(\mathbb{P}^2)^\vee$. This means the curves lie in the double cover of $(\mathbb{P}^2)^\vee$ branched over a sextic, i.e., a K3 surface.



Fibrations by products of elliptic curves

Thm (Kamenova): Let $X \rightarrow \mathbb{P}^2$ be a Lagrangian fibration with

- X smooth,
- general fibre a product of two elliptic curves,
- “generic” singular fibres,
- and a global section.

Then X is birational to $\text{Hilb}^2 S$ of an elliptic K3 surface S .

Thm (Debarre-Huybrechts-Macri-Voisin): Let $X \rightarrow \mathbb{P}^2$ be a (numerical!) Lagrangian fibration with X smooth and a divisor $Y \subset X$ inducing a principal polarization on a general fibre. Then X is a deformation of $\text{Hilb}^2 S$.

Rmk: These results cover examples 1a and 1b in four dimensions. Next consider example 3a with $(1, 2)$ -polarized fibres.

Fibrations by $(1, 2)$ -polarized fibres

If A is $(1, 2)$ -polarized then A^\vee is too. Let $C \subset A^\vee$ be a polarization. Then C is genus three, and pull-back gives

$$A = \text{Pic}^0 A^\vee \longrightarrow \text{Jac}^0 C \longrightarrow E,$$

i.e., A is the Prym variety of a double cover $C \rightarrow E$.

Thm (Qin-S-): Let $\mathcal{C}/\mathcal{E}/\mathbb{P}^2$ be a flat family of double covers of reduced Gorenstein curves of genus three and one, respectively, such that $X = \overline{\text{Prym}}(\mathcal{C}/\mathcal{E})$ is a Lagrangian fibration. Then $X \rightarrow \mathbb{P}^2$ must be a Markushevich-Tikhomirov system

Thus the elliptic curves \mathcal{E} must lie in a degree two del Pezzo and the genus three curves \mathcal{C} must lie in its K3 double cover.

Fibrations by (1, 2)-polarized fibres

Proof: $f : \mathcal{C} \rightarrow \mathcal{E}$ is branched over a divisor of degree four on each fibre. Thus there is a line bundle \mathcal{L} of degree two on each fibre with

$$f_*\mathcal{O}_{\mathcal{C}} = \mathcal{O}_{\mathcal{E}} \oplus \mathcal{L}^{\vee}.$$

Applying $R^1\pi_*$ gives (π always denotes projection to \mathbb{P}^2)

$$R^1\pi_*\mathcal{O}_{\mathcal{C}} = R^1\pi_*\mathcal{O}_{\mathcal{E}} \oplus R^1\pi_*\mathcal{L}^{\vee}.$$

On the other hand, $\text{Jac}^0\mathcal{C} \sim \text{Jac}^0E \times \text{Prym}(C/E)$ implies

$$H^1(C, \mathcal{O}_C) = H^1(E, \mathcal{O}_E) \oplus H^1(X_t, \mathcal{O}_{X_t}),$$

$$R^1\pi_*\mathcal{O}_{\mathcal{C}} = R^1\pi_*\mathcal{O}_{\mathcal{E}} \oplus R^1\pi_*\mathcal{O}_X.$$

Fibrations by $(1, 2)$ -polarized fibres

Therefore

$$R^1\pi_*\mathcal{L}^\vee \cong R^1\pi_*\mathcal{O}_X \cong \Omega_{\mathbb{P}^2}^1,$$

and we have a double cover

$$h : \mathcal{E} \longrightarrow \mathbb{P}(R^1\pi_*\mathcal{L}^\vee) = \mathbb{P}(\Omega_{\mathbb{P}^2}^1) \subset \mathbb{P}^2 \times (\mathbb{P}^2)^\vee$$

branched over the zero locus in $\mathbb{P}(\Omega_{\mathbb{P}^2}^1)$ of a section of

$$\mathcal{O}_{\mathbb{P}(\Omega^1)}(4) \otimes \pi^*\mathcal{O}_{\mathbb{P}^2}(2d) = \mathcal{O}(2d + 4, 4)|_{\mathbb{P}(\Omega^1)}.$$

Recall $\mathcal{C} \rightarrow \mathcal{E}$ is branched over the zero locus in \mathcal{E} of a section of

$$\mathcal{L}^2 \cong h^*(\mathcal{O}_{\mathbb{P}(\Omega^1)}(2) \otimes \pi^*\mathcal{O}_{\mathbb{P}^2}(e)) = h^*\mathcal{O}(e + 2, 2)|_{\mathbb{P}(\Omega^1)}.$$

Now $R^1\pi_*\mathcal{L}^\vee = \Omega_{\mathbb{P}^2}^1$ determines $d = -2$ and $e = -2$.

Fibrations by $(1, 2)$ -polarized fibres

So

$$\mathcal{O}(2d + 4, 4)|_{\mathbb{P}(\Omega^1)} = \mathcal{O}(0, 4)|_{\mathbb{P}(\Omega^1)},$$

$$h^* \mathcal{O}(e + 2, 2)|_{\mathbb{P}(\Omega^1)} = h^* \mathcal{O}(0, 2)|_{\mathbb{P}(\Omega^1)}$$

are pulled back from $(\mathbb{P}^2)^\vee$.

This means the elliptic curves \mathcal{E} lie in the double cover of $(\mathbb{P}^2)^\vee$ branched over a quartic, i.e., a degree two del Pezzo surface T .

Rmk: If $g : T \rightarrow (\mathbb{P}^2)^\vee$ then

$$K_T \cong g^*(\mathcal{O}(-3) \otimes \mathcal{O}(2)) = g^*\mathcal{O}(-1).$$

Moreover, the genus three curves \mathcal{C} lie in the double cover of T branched over the pull-back of a conic $\cong K_T^{-2}$, i.e., a K3 surface.



Fibrations by Jacobians in six dimensions

Thm (S-): Let \mathcal{C}/\mathbb{P}^3 be a flat family of reduced curves of genus 3 such that $X = \overline{\text{Jac}}^d(\mathcal{C}/\mathbb{P}^3)$ is a Lagrangian fibration. If the curves

- are irreducible Gorenstein non-hyperelliptic, or
- are canonically positive 2-connected hyperelliptic,

then X/\mathbb{P}^3 must be a Beauville-Mukai integrable system.

Rmk: The general principally polarized abelian threefold is the Jacobian of a (non-hyperelliptic) curve of genus three.

Qu: If the general fibre of X/\mathbb{P}^3 is the product $E_1 \times E_2 \times E_3$ of elliptic curves, must X be $\text{Hilb}^3 S$ of an elliptic K3 S ?

Fibrations by Jacobians in six dimensions

Proof (in the non-hyperelliptic case): The relative canonical embedding gives

$$\mathcal{C} \rightarrow \mathbb{P}(R^1\pi_*\mathcal{O}_{\mathcal{C}}) = \mathbb{P}(R^1\pi_*\mathcal{O}_X) = \mathbb{P}(\Omega_{\mathbb{P}^3}^1) \subset \mathbb{P}^3 \times (\mathbb{P}^3)^\vee.$$

Indeed \mathcal{C} is the zero locus in $\mathbb{P}(\Omega_{\mathbb{P}^3}^1)$ of a section of

$$\mathcal{O}_{\mathbb{P}(\Omega^1)}(4) \otimes \pi^*\mathcal{O}_{\mathbb{P}^3}(k) = \mathcal{O}(k+4, 4)|_{\mathbb{P}(\Omega^1)}.$$

Now $R^1\pi_*\mathcal{O}_{\mathcal{C}} = \Omega_{\mathbb{P}^3}^1$ determines $k = -4$, so

$$\mathcal{O}(k+4, 4)|_{\mathbb{P}(\Omega^1)} = \mathcal{O}(0, 4)|_{\mathbb{P}(\Omega^1)}$$

is pulled back from $(\mathbb{P}^3)^\vee$. This means the curves are hyperplane sections of a quartic K3 surface in $(\mathbb{P}^3)^\vee$.



An invariant of Lagrangian fibrations

Thm (Wieneck): In a family of Lagrangian fibrations the polarization type of the fibres is constant.

For fibrations on $\text{Hilb}^n K3$ the polarization is *principal*

$$(d_1, d_2, \dots, d_n) = (1, 1, \dots, 1).$$

For fibrations on $K_n(A)$ the polarization is of type

$$(d_1, d_2, \dots, d_{n-2}, d_{n-1}, d_n) = (1, 1, \dots, 1, d_{n-1}, d_n)$$

with $d_{n-1}d_n = n + 1$.

Thm (Markman): If X is a general deformation of $\text{Hilb}^n K3$ admitting a Lagrangian fibration then it is birational to a *Tate-Shafarevich twist* of Beauville-Mukai system.

Finiteness

Thm (S-): Fixing $d_1 | \dots | d_n$, there are finitely many Lagrangian fibrations up to deformation with

- polarization type (d_1, \dots, d_n) ,
- a global section,
- maximally varying fibres,
- semistable singular fibres in codimension one.

Rmk: van Geemen-Voisin and Bakker proved that Lagrangian fibrations are maximally varying or *isotrivial*.

Rmk: Using a theorem of Charles, Kamenova showed that it is enough to assume there is a fibration with a fixed polarization type. (See also Debarre-Huybrechts-Macri-Voisin.)

Thus, we want to bound the polarization type.

Restrictions on polarization type

Thm (S-): Let $X \rightarrow \mathbb{P}^2$ be a Lagrangian fibration (with X smooth!) and let $Y \in H^2(X, \mathbb{Z})$ restrict to a polarization of type (d_1, d_2) on each smooth fibre. Then $d_1 d_2$ can take only the following twenty values up to a square:

$$1, 2, 3, 5, 7, 10, 15, 61, 62, 241, 246, 247, \\ 249, 251, 253, 254, 255, 257, 258, 259$$

Example: Polarization types $(1, 6)$ and $(1, 11)$ are *not* possible. Debarre-Huybrechts-Macri-Voisin showed that $(1, 2)$, $(1, 5)$, and $(1, 7)$ are also *not* possible (for X smooth!).

Rmk: In higher dimensions, the overall degree $d_1 \cdots d_n$ of the fibres can take only finitely many values up to an n^{th} power.

Restrictions on polarization type

Proof: Let L be the pullback of a hyperplane from \mathbb{P}^2 . Then

$$\left(\int_X Y^2 L^2 \right) \left(\int_X c_2(\sigma\bar{\sigma}) \right)^2 = \left(\int_X (\sigma\bar{\sigma})^2 \right) \left(\int_X c_2 YL \right)^2$$

where $\int_X Y^2 L^2 = 2!d_1d_2$ and Hitchin-S- formula gives

$$\frac{\left(\int_X c_2(\sigma\bar{\sigma}) \right)^2}{\int_X (\sigma\bar{\sigma})^2} = \frac{24^2(2!)^2}{2^2} \sqrt{\hat{A}[X]}.$$

Thus

$$1152d_1d_2\sqrt{\hat{A}[X]} = \left(\int_X c_2 YL \right)^2 \text{ is a square.}$$

Guan showed that $\sqrt{\hat{A}[X]}$ takes finitely many values.



Summary of examples in four dimensions

Example	Polarization type
Beauville-Mukai system	(1, 1)
Hilb ² <i>S</i> of an elliptic K3 <i>S</i>	(1, 1)
Arbarello-Saccà-Ferretti system	(1, 1)
Isotrivial system on Hilb ² <i>S</i> / $\mathbb{Z}_2^{\oplus 2}$	(1, 1)
Markushevich-Tikhomirov system	(1, 2)
Isotrivial system on Hilb ² <i>S</i> / \mathbb{Z}_2	(1, 2)
Debarre system for $A^{1,3}$	(1, 3)
Isotrivial system on $K_2(E \times F)$	(1, 3)

Summary of examples in six dimensions

Example	Polarization type
Beauville-Mukai system	(1, 1, 1)
Hilb ³ S of an elliptic K3 S	(1, 1, 1)
Arbarello-Saccà-Ferretti system	(1, 1, 1)
Matteini system	(1, 1, 2)
S-Shen system	(1, 2, 2)
Debarre system for $A^{1,4}$	(1, 1, 4)
Isotrivial system on $K_3(A)$	(1, 1, 4)
Debarre system for $A^{2,2}$	(1, 2, 2)
O'Grady 6 on $A^{2,2}$	(1, 2, 2)

Summary of examples in six dimensions with duals

Example	Type	Type	Dual
Beauville-Mukai	(1, 1, 1)	(1, 1, 1)	Beauville-Mukai
Hilb ³ of elliptic K3	(1, 1, 1)	(1, 1, 1)	Hilb ³ of elliptic K3
ASF system	(1, 1, 1)	(1, 1, 1)	ASF system
Matteini system	(1, 1, 2)		
S-Shen system	(1, 2, 2)	(1, 1, 2)	degenerate Matteini?
Debarre for $A^{1,4}$	(1, 1, 4)	(1, 4, 4)	Kim/S-
Isotrivial on $K_3(A)$	(1, 1, 4)	(1, 4, 4)	Kim/S-
Debarre for $A^{2,2}$	(1, 2, 2)	(1, 1, 2)	Kim/S-
O'Grady 6 on $A^{2,2}$	(1, 2, 2)	(1, 1, 2)	Kim/S-